arXiv:1609.07385v1 [math-ph] 23 Sep 2016

Exact solution of the relativistic quantum Toda chain

Xin Zhang[a], Junpeng Cao[a,b,c], Wen-Li Yang[d,e,f1], Kangjie Shi[d,e] and Yupeng Wang[a,b,c2]

[a]Beijing National Laboratory for Condensed Matter Physics, Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China

[b]Collaborative Innovation Center of Quantum Matter, Beijing, China

[c]School of Physical Sciences, University of Chinese Academy of Sciences, Beijing, China

[d]Institute of Modern Physics, Northwest University, Xian 710069, China

[e]Shaanxi Key Laboratory for Theoretical Physics Frontiers, Xian 710069, China

[f]Beijing Center for Mathematics and Information Interdisciplinary Sciences, Beijing, 100048, China

Abstract

The relativistic quantum Toda chain model is studied with the generalized algebraic Bethe Ansatz method. By employing a set of local gauge transformations, proper local vacuum states can be obtained for this model. The exact spectrum and eigenstates of the model are thus constructed simultaneously.

PACS: 02.30.Ik, 03.65.Vf, 02.10.De
Keywords: Toda chain, Generalized Bethe Ansatz, $T-Q$ relation

[1]Corresponding author: wlyang@nwu.edu.cn
[2]Corresponding author: yupeng@iphy.ac.cn

1 Introduction

The Toda chain model [1, 2] is one of the most fascinating integrable models [3–5], which plays an important role in theoretical physics and mathematics such as in Yang-Mills field theory [6] and group theory [7]. In the past several decades, considerable attentions have been paid on both the classical Toda chain [5, 8, 9] and the quantum Toda chain [10–15] because of its unique property: an infinite dimensional model without $U(1)$ symmetry. This unique property makes it almost impossible to derive its spectrum with the conventional Bethe Ansatz methods such as the coordinate Bethe Ansatz and the algebraic Bethe Ansatz since the absence of an obvious reference state. To overcome this difficulty, several remarkable approaches have then been developed such as Gutzwiller's Ansatz [10] for few-body systems, asymptotic Bethe Ansatz [11] and functional Bethe Ansatz [12–14]. An interesting issue is that the relativistic Toda chain model [16–19], which is tightly related to the Seiberg-Witten theory [6] and the Calabi-Yau manifold [20–22], is also integrable. However, its spectrum has not been derived exactly yet.

In this paper, we study the relativistic quantum Toda chain [17, 18] described by the Hamiltonian

$$H=\sum_{n=1}^{N}\cos(2\eta\hat{p}_n)+\sum_{n=1}^{N}g^2\cos(\eta\hat{p}_n+\eta\hat{p}_{n+1})\,e^{x_{n+1}-x_n}, \tag{1.1}$$

where η and g are two generic coupling constants; x_n and $\hat{p}_n$ denote the coordinate and momentum operators on site n which satisfy the commutation relation: $[x_n,\hat{p}_m]=i\delta_{m,n}$. Notice that the periodic boundary condition $x_{N+1}=x_1$ and $\hat{p}_{N+1}=\hat{p}_1$ is imposed in (1.1). The non-relativistic Toda chain Hamiltonian $\mathcal{H}$ can be readily obtained by taking the limit $\eta\to 0$ and $g=i\sqrt{2c}\eta\to 0$ with c a constant. In this limit the Hamiltonian (1.1) has the expansion

$$H=N-2\eta^2\,\mathcal{H}+\cdots,$$

and the resulting $\mathcal{H}$ is exactly the Hamiltonian of the non-relativistic Toda chain [1, 2] given by

$$\mathcal{H}=\sum_{n=1}^{N}\hat{p}_n^2+c\sum_{n=1}^{N}e^{x_{n+1}-x_n}. \tag{1.2}$$

Here we shall adopt the generalized Bethe Ansatz method used in [23, 24] to construct the exact spectrum of Hamiltonian (1.1).

The paper is organized as follows. In Section 2, we briefly review the integrability of the relativistic quantum Toda chain by constructing the associated commuting transfer matrix. A set of gauge transformations and the corresponding local vacuum states are given in Section 3. Some useful commutation relations among the matrix elements of the gauged monodromy matrix are also presented. The exact spectrum given in terms of a homogeneous T-Q relation and the associated Bethe Ansatz equations (BAEs) and eigenstates of the transfer matrix are derived in Section 4. Concluding remarks are given in Section 5. Some useful formulas and another set of gauge transformations are shown in Appendix A & B, respectively.

2 Integrability

Let $\mathbf{V}$ denote the local Hilbert space. The Lax operator of the relativistic quantum Toda chain $L_n(u) \in \mathrm{End}(\mathbf{C}^2 \otimes \mathbf{V})$ with a generic spectral parameter u is defined as [17, 18]

$$L_n(u) = \begin{pmatrix} e^{u-i\eta\hat{p}_n} - e^{-u+i\eta\hat{p}_n} & -ge^{x_n} \\ ge^{-x_n} & 0 \end{pmatrix}, \quad n = 1, \cdots, N, \tag{2.1}$$

which satisfies the Yang-Baxter relation

$$R(u-v)(L_n(u) \otimes I)(I \otimes L_n(v)) = (I \otimes L_n(v))(L_n(u) \otimes I)R(u-v), \quad n = 1, \cdots, N, \tag{2.2}$$

where the R-matrix $R(u) \in \mathrm{End}(\mathbf{C}^2 \otimes \mathbf{C}^2)$ reads

$$R(u) = \begin{pmatrix} \sinh(u+\eta) & 0 & 0 & 0 \\ 0 & \sinh u & \sinh \eta & 0 \\ 0 & \sinh \eta & \sinh u & 0 \\ 0 & 0 & 0 & \sinh(u+\eta) \end{pmatrix}. \tag{2.3}$$

The monodromy matrix of the model is constructed as

$$T(u) = L_N(u) \cdots L_1(u) = \begin{pmatrix} A(u) & B(u) \\ C(u) & D(u) \end{pmatrix}. \tag{2.4}$$

With the help of Eq.(2.2), we conclude that the monodromy matrix (2.4) also satisfies the Yang-Baxter equation

$$R_{1,2}(u_1 - u_2)T_1(u_1)T_2(u_2) = T_2(u_2)T_1(u_1)R_{1,2}(u_1 - u_2). \tag{2.5}$$

Here and below we adopt the standard notaitons: for any matrix $A \in \mathrm{End}(\mathbf{V})$, A_n is an embedding operator in the tensor space $\mathbf{V} \otimes \mathbf{V} \otimes \cdots$, which acts as A on site n and an

identity on the other factor spaces; $R_{i,j}(u)$ is an embedding operator in the tensor space, which acts as identity on the factor space except for the i-th and j-th ones.

As usual, the transfer matrix $t(u)$ of the present model is given by

$$t(u) = \mathrm{tr}T(u) = A(u) + D(u) = \sum_{j=0}^{N} t_{N-2j} e^{(N-2j)u}.$$

Eq.(2.5) leads to that the transfer matrix $t(u)$ forms a mutually commuting family [25–27], i.e., $[t(u), t(v)] = 0$. Therefore, $t(u)$ serves as a generating functional of conserved quantities. The Hamiltonian of the relativistic quantum Toda chain (1.1) can be obtained through the resulting conserved quantities by [18]

$$H = -\frac{1}{2}\left(t_{N-2}t_N^{-1} + t_{2-N}t_{-N}^{-1}\right). \tag{2.6}$$

3 Gauge transformation and local vaccum

For generic complex η with $Re(\eta) > 0$, to find a proper vacuum state which allows us to perform the algebraic Bethe Ansatz, we introduce the following gauge matrices

$$M_k(u) = \left(\begin{array}{cc} X_k(u), & Y_k(u) \end{array}\right) = \begin{pmatrix} \frac{e^{-u-k\eta}}{\sinh(k\eta)} & e^{-u+k\eta} \\ \frac{1}{\sinh(k\eta)} & 1 \end{pmatrix},$$

$$M_k^{-1}(u) = \begin{pmatrix} \overline{Y}_k(u) \\ \overline{X}_k(u) \end{pmatrix} = \frac{e^u}{2} \begin{pmatrix} -1 & e^{-u+k\eta} \\ \frac{1}{\sinh(k\eta)} & -\frac{e^{-u-k\eta}}{\sinh(k\eta)} \end{pmatrix}, \tag{3.1}$$

where k is a free complex parameter. The matrices $L_n(u)$ and $T(u)$ under the gauge transformations behave as

$$\overline{L}_{j,k}^{(n)}(u) = M_j^{-1}(u) L_n(u) M_k(u) = \begin{pmatrix} \overline{A}_{j,k}^{(n)}(u) & \overline{B}_{j,k}^{(n)}(u) \\ \overline{C}_{j,k}^{(n)}(u) & \overline{D}_{j,k}^{(n)}(u) \end{pmatrix}, \quad n = 1, \cdots, N, \tag{3.2}$$

$$\overline{T}_{j,k}(u) = M_j^{-1}(u) T(u) M_k(u) = \begin{pmatrix} \overline{A}_{j,k}(u) & \overline{B}_{j,k}(u) \\ \overline{C}_{j,k}(u) & \overline{D}_{j,k}(u) \end{pmatrix}. \tag{3.3}$$

Now we define a local wave function on site n as

$$|\alpha; n\rangle = e^{-\frac{1}{2\eta}(x_n - \alpha\eta)^2 + \beta_n x_n}, \quad n = 1, \cdots, N, \tag{3.4}$$

with α a free parameter and

$$\beta_n = -n - \frac{1}{2} - \frac{(2n+1)\ln g + in\pi}{\eta}. \tag{3.5}$$

As we consider the $Re(\eta) > 0$ case, the wave function (3.4) is convergent in the whole real x_n line. When $Re(\eta) < 0$ (the Hamiltonian is in fact an even function of η), we can use another set of gauge transformation listed in Appendix B to obtain the exact solutions of this model. The $Re(\eta) = 0$ case will be discussed in Section 4. With the help of the definitions (3.2) and (3.4), we can easily check that

$$\overline{B}^{(n)}_{\alpha_{n+1},\alpha_n}(u)|\alpha; n\rangle = 0, \tag{3.6}$$

$$\overline{A}^{(n)}_{\alpha_{n+1},\alpha_n}(u)|\alpha; n\rangle = ge^{\frac{\eta}{2}}e^{u-n\delta\eta}|\alpha+1; n\rangle, \tag{3.7}$$

$$\overline{D}^{(n)}_{\alpha_{n+1},\alpha_n}(u)|\alpha; n\rangle = ge^{\frac{\eta}{2}}e^{-u+n\delta\eta}|\alpha-1; n\rangle, \tag{3.8}$$

where

$$\delta = -1 - \frac{2\ln g + i\pi}{\eta}, \tag{3.9}$$

$$\alpha_n = \alpha + n\delta, \quad n = 1, \cdots, N+1. \tag{3.10}$$

Let us define the global vacuum state

$$|\alpha\rangle = \otimes_{n=1}^{N}|\alpha; n\rangle. \tag{3.11}$$

Acting the elements of the gauge transformed matrix (3.3) on the state (3.11), we readily have

$$\overline{B}_{\alpha_{N+1},\alpha_1}(u)|\alpha\rangle = 0, \tag{3.12}$$

$$\overline{A}_{\alpha_{N+1},\alpha_1}(u)|\alpha\rangle = a(u)|\alpha+1\rangle, \tag{3.13}$$

$$\overline{D}_{\alpha_{N+1},\alpha_1}(u)|\alpha\rangle = d(u)|\alpha-1\rangle, \tag{3.14}$$

where

$$a(u) = g^N e^{\frac{N\eta}{2}}e^{Nu-\frac{N(N+1)\delta\eta}{2}}, \qquad d(u) = g^N e^{\frac{N\eta}{2}}e^{-Nu+\frac{N(N+1)\delta\eta}{2}}. \tag{3.15}$$

Therefore $|\alpha\rangle$ serves as a reference state in the sense of Ref. [23].

In addition, with the help of the Yang-Baxter relation (2.5), we can derive the commutation relations

$$\overline{C}_{m',m}(u_1)\overline{C}_{m'+1,m-1}(u_2) = \overline{C}_{m',m}(u_2)\overline{C}_{m'+1,m-1}(u_1), \tag{3.16}$$

$$\overline{A}_{m',m}(u_1)\overline{C}_{m'+1,m-1}(u_2)=\frac{\sinh(u_1-u_2+\eta)}{\sinh(u_1-u_2)}\overline{C}_{m'+2,m}(u_2)\overline{A}_{m'+1,m-1}(u_1)$$
$$-\frac{\sinh\eta\sinh((m'+1)\eta-u_1+u_2)}{\sinh(u_1-u_2)\sinh(m'+1)\eta}\overline{C}_{m'+2,m}(u_1)\overline{A}_{m'+1,m-1}(u_2), \tag{3.17}$$
$$\overline{D}_{m',m}(u_1)\overline{C}_{m'+1,m-1}(u_2)=\frac{\sinh(u_1-u_2-\eta)}{\sinh(u_1-u_2)}\overline{C}_{m',m-2}(u_2)\overline{D}_{m'+1,m-1}(u_1)$$
$$+\frac{\sinh\eta\sinh((m-1)\eta-u_1+u_2)}{\sinh(u_1-u_2)\sinh(m-1)\eta}\overline{C}_{m',m-2}(u_1)\overline{D}_{m'+1,m-1}(u_2), \tag{3.18}$$

which together with the global vacuum state (3.11) allow us to perform the generalized algebraic Bethe Ansatz [23, 24]. The proof of the above commutation relations is given in Appendix A.

4 Generalized Bethe Ansatz

Let us consider first the case of a special sequence of η taking values

$$\eta=\frac{i\pi(2q-N)}{N+2M}-\frac{2N\ln g}{N+2M},\quad M=0,1,2,\cdots\quad\text{and}\quad q\in\mathbb{Z}, \tag{4.1}$$

and $|g|<1$ (for $|g|>1$ case we can use the gauge transformation introduced in Appendix B). We remark that when the parameter η takes the above discrete values the identity: $e^{2M\eta}=e^{N\delta\eta}$ holds, which allows us to introduce the Bethe-type state

$$|u_1,\cdots,u_M;\alpha\rangle=\left\{\prod_{j=1}^{M}\overline{C}_{k_\alpha+j,k_\alpha-j}(u_j)\right\}|\alpha\rangle, \tag{4.2}$$

where

$$k_\alpha=\alpha+\delta+M. \tag{4.3}$$

Acting $\overline{A}_{k_\alpha,k_\alpha}(u)$ and $\overline{D}_{k_\alpha,k_\alpha}(u)$ on the state $|u_1,\cdots,u_M;\alpha\rangle$ and using the commutation relations (3.16)-(3.18), we have

$$\overline{A}_{k_\alpha,k_\alpha}(u)|u_1,\cdots,u_M;\alpha\rangle=a(u;\{u_l\})|u_1,\cdots,u_M;\alpha+1\rangle$$
$$+\sum_{j=1}^{M}a_j(u;k_\alpha;\{u_l\})|u_1,\cdots,u_{j-1},u,u_{j+1},\cdots,u_M;\alpha+1\rangle, \tag{4.4}$$
$$\overline{D}_{k_\alpha,k_\alpha}(u)|u_1,\cdots,u_M;\alpha\rangle=d(u;\{u_l\})|u_1,\cdots,u_M;\alpha-1\rangle$$
$$+\sum_{j=1}^{M}d_j(u;k_\alpha;\{u_l\})|u_1,\cdots,u_{j-1},u,u_{j+1},\cdots,u_M;\alpha-1\rangle, \tag{4.5}$$

where

$$
\begin{aligned}
a(u;\{u_l\}) &= a(u)\prod_{j=1}^{M}\frac{\sinh(u-u_j+\eta)}{\sinh(u-u_j)},\\
d(u;\{u_l\}) &= d(u)\prod_{j=1}^{M}\frac{\sinh(u-u_j-\eta)}{\sinh(u-u_j)},\\
a_j(u;k_\alpha;\{u_l\}) &= -a(u_j)\frac{\sinh\eta\sinh((k_\alpha+1)\eta-u+u_j)}{\sinh(u-u_j)\sinh((k_\alpha+1)\eta)}\prod_{l\neq j}^{M}\frac{\sinh(u_j-u_l+\eta)}{\sinh(u_j-u_l)},\\
d_j(u;k_\alpha;\{u_l\}) &= d(u_j)\frac{\sinh\eta\sinh((k_\alpha-1)\eta-u+u_j)}{\sinh(u-u_j)\sinh((k_\alpha-1)\eta)}\prod_{l\neq j}^{M}\frac{\sinh(u_j-u_l-\eta)}{\sinh(u_j+u_l)}.
\end{aligned}
$$

Assume that an eigenstate of the transfer matrix $t(u)$ takes the following form

$$
|\lambda_1,\cdots,\lambda_M;\bar{\alpha}\rangle\rangle=\sum_{n\in\mathbb{Z}}e^{i(\bar{\alpha}+n)\phi}|\lambda_1,\cdots,\lambda_M;\bar{\alpha}+n\rangle, \tag{4.6}
$$

where ϕ is a complex parameter and $\bar{\alpha}\neq-\delta+j+\frac{ik\pi}{\eta}$, $j,k\in\mathbb{Z}$. The definition of transfer matrix $t(u)$ implies that

$$
t(u)=\mathrm{tr}T(u)=\mathrm{tr}\overline{T}_{k,k}(u)=\overline{A}_{k,k}(u)+\overline{D}_{k,k}(u). \tag{4.7}
$$

Applying $t(u)$ on the eigenstate (4.6) and using the relations (4.4) and (4.5), we have

$$
\begin{aligned}
t(u)|\lambda_1,\cdots,\lambda_M;\bar{\alpha}\rangle\rangle=\Lambda(u)|\lambda_1,\cdots,\lambda_M;\bar{\alpha}\rangle\rangle+\sum_{j}^{M}\Lambda_j(u)\Bigg\{\frac{\sinh\eta}{\sinh(u-\lambda_j)}\sum_{n\in\mathbb{Z}}e^{i(\bar{\alpha}+n)\phi}\\
\times\frac{\sinh(u-\lambda_j-(k_{\bar{\alpha}}+n)\eta)}{\sinh((k_{\bar{\alpha}}+n)\eta)}|\lambda_1,\cdots,\lambda_{j-1},u,\lambda_{j+1},\cdots,\lambda_M;\bar{\alpha}+n\rangle\Bigg\},
\end{aligned} \tag{4.8}
$$

where $k_{\bar{\alpha}}=\bar{\alpha}+\delta+M$. The function $\Lambda(u)$ in (4.8) identifies the eigenvalue term

$$
\Lambda(u)=e^{-i\phi}a(u)\prod_{j=1}^{M}\frac{\sinh(u-\lambda_j+\eta)}{\sinh(u-\lambda_j)}+e^{i\phi}d(u)\prod_{j=1}^{M}\frac{\sinh(u-\lambda_j-\eta)}{\sinh(u-\lambda_j)}, \tag{4.9}
$$

and the coefficients $\{\Lambda_j(u)|j=1,\cdots,M\}$ of the unwanted terms read

$$
\Lambda_j(u)=e^{-i\phi}a(\lambda_j)\prod_{l\neq j}^{M}\frac{\sinh(\lambda_j-\lambda_l+\eta)}{\sinh(\lambda_j-\lambda_l)}-e^{i\phi}d(\lambda_j)\prod_{l\neq j}^{M}\frac{\sinh(\lambda_j-\lambda_l-\eta)}{\sinh(\lambda_j-\lambda_l)}. \tag{4.10}
$$

To ensure the state (4.6) to be an eigenstate of the transfer matrix, we should put $\Lambda_j(u)=0$ which gives rise to the associated BAEs

$$e^{2i\phi}\frac{d(\lambda_j)}{a(\lambda_j)}=\prod_{l\neq j}^{M}\frac{\sinh(\lambda_j-\lambda_l+\eta)}{\sinh(\lambda_j-\lambda_l-\eta)},\quad j=1,\cdots,M. \tag{4.11}$$

To determine the parameter ϕ, we should note that the transfer matrix $t(u)$ possesses the asymptotic behavior

$$\lim_{u\to\pm\infty}t(u)=t_{\pm N}e^{\pm Nu}+\cdots=(\pm 1)^N e^{\mp i\eta\sum_n\hat{p}_n}e^{\pm Nu}+\cdots.$$

The commutation relation $[t(u),t_n]=0$ implies that

$$\lim_{u\to\pm\infty}\Lambda(u)=(\pm 1)^N e^{\mp i\eta K}e^{\pm Nu}+\cdots,$$

where $K\equiv\sum_n p_n$ is the total momentum of the system. The above asymptotic behavior allows us to fix the parameter ϕ as

$$e^{i\phi}=g^N e^{-NM\eta}e^{\frac{N\eta}{2}+i\eta K},\quad \text{or}\quad e^{-i\phi}=(-1)^N g^N e^{(N+2)M\eta}e^{\frac{N\eta}{2}-i\eta K}, \tag{4.12}$$

which is self-consistent with each other for η values given in equation (4.1). The $T-Q$ relation given by (4.9) can be rewritten as

$$\begin{aligned}\Lambda(u) &= (ig)^N e^{\frac{N\eta}{2}}e^{Nu-i\eta K}\prod_{j=1}^{M}\frac{Q(u+\eta)}{Q(u)}+(-ig)^N e^{\frac{N\eta}{2}}e^{-Nu+i\eta K}\prod_{j=1}^{M}\frac{Q(u-\eta)}{Q(u)}, \\ Q(u) &= \prod_{j=1}^{M}\sinh(u-\lambda_j),\end{aligned} \tag{4.13}$$

where $K\in\mathbb{R}$ is the total momentum of the system and the BAEs (4.11) become

$$e^{-2N\lambda_j+2i\eta K}=(-1)^N\prod_{l\neq j}^{M}\frac{\sinh(\lambda_j-\lambda_l+\eta)}{\sinh(\lambda_j-\lambda_l-\eta)},\qquad j=1,\cdots,M. \tag{4.14}$$

The $T-Q$ relation (4.13) is the eigenvalue of the transfer matrix with the corresponding eigenstste $|\lambda_1,\cdots,\lambda_M;\bar{\alpha}\rangle\rangle$ given by (4.6) and (4.12), provided that the parameters $\{\lambda_j|j=1,\cdots,M\}$ satisfy the BAEs (4.14).

With the help of the definition (2.6), we can easily obtain the eigenvalue of the Hamiltonian in terms of the Bethe roots as

$$E=(e^{-2\eta}-1)\sum_{j=1}^{M}\cosh(2\lambda_j). \tag{4.15}$$

Now let us consider the generic imaginary η case which together with real g^2 defines the physically meaningful relativistic quantum Toda chain. Keeping N fixed and taking $M, q \to \infty$ but with $q/M \to$ finite, the η values in (4.1) become dense and tend to generic imaginary values. In such a limit, we conclude that the Q-function is an infinite product for $|g| \neq 1$, quite similar to that of the non-relativistic Toda chain obtained by Sklyanin [12] due to the fact that the model is infinite dimensional and without $U(1)$ symmetry. The resulting eigenvalues and the BAEs read

$$\begin{aligned}\Lambda(u) &= (ig)^N e^{\frac{N\eta}{2}} e^{Nu - i\eta K} \prod_{j=1}^{\infty} \frac{\sinh(u-\lambda_j+\eta)}{\sinh(u-\lambda_j)} \\ &\quad + (-ig)^N e^{\frac{N\eta}{2}} e^{-Nu + i\eta K} \prod_{j=1}^{\infty} \frac{\sinh(u-\lambda_j-\eta)}{\sinh(u-\lambda_j)}, \end{aligned} \tag{4.16}$$

and

$$e^{-2N\lambda_j + 2i\eta K} = (-1)^N \prod_{l\neq j}^{\infty} \frac{\sinh(\lambda_j - \lambda_l + \eta)}{\sinh(\lambda_j - \lambda_l - \eta)}, \qquad j = 1, \cdots, M, \tag{4.17}$$

with the constraint condition

$$\lim_{u\to\pm\infty} \prod_{j=1}^{\infty} \frac{\sinh(u-\lambda_j \pm \eta)}{\sinh(u-\lambda_j)} = (ig)^{-N} e^{-\frac{N\eta}{2}}. \tag{4.18}$$

A very special case is that of $|g| = 1$. In this case, the wave function is somehow subtle but the spectrum can be extrapolated from (4.13)-(4.14) by taking the limit $Re(\eta) \to 0^+$. We remark that with the cyclic representation of the Weyl algebra [16, 19] (replacing $e^{i\eta\hat{p}_n}$ and e^{x_n} in the present paper by $e^{\eta\hat{p}_n}$ and e^{ix_n}), the model at roots of unit is reduced to a special case of the quantum τ_2-model (finite dimensional) and can be solved exactly via the off-diagonal Bethe Ansatz [29].

5 Conclusion

By employing a set of gauge transformations (3.1), we have obtained the proper vacuum state (3.11) which allows us to use the generalized algebraic Bethe ansatz [23,24] to construct the eigenvalues (4.15), the corresponding eigenstates (4.6) and the associated BAEs (4.14) of the relativistic quantum Toda chain when the parameter η takes some discrete values (4.1). Keeping N fixed and taking $M, q \to \infty$ but with $q/M \to$ finite, the η values in (4.1) become dense and tend to generic imaginary values. Thus the resulting results give the exact solutions of the relativistic quantum Toda chain.

Acknowledgments

The financial supports from the National Natural Science Foundation of China (Grant Nos. 11375141, 11374334, 11434013, 11425522), BCMIIS, the National Program for Basic Research of MOST (Grant No. 2016YFA0300603) and the Strategic Priority Research Program of the Chinese Academy of Sciences are gratefully acknowledged.

Appendix A: Commutation relations

From the definition of the R-matrix (2.3) and the gauge transformation (3.1), we can obtain the following useful commutation relations [24, 28]

$$R_{1,2}(u_1-u_2)X^1_m(u_1)X^2_{m-1}(u_2)=\sinh(u_1-u_2+\eta)X^2_m(u_2)X^1_{m-1}(u_1), \tag{A.1}$$

$$R_{1,2}(u_1-u_2)Y^1_m(u_1)Y^2_{m+1}(u_2)=\sinh(u_1-u_2+\eta)Y^2_m(u_2)Y^1_{m+1}(u_1), \tag{A.2}$$

$$\begin{aligned}R_{1,2}(u_1-u_2)X^1_{m-1}(u_1)Y^2_m(u_2)&=\sinh(u_1-u_2)Y^2_{m+1}(u_2)X^1_m(u_1)\\&+\frac{\sinh\eta\sinh(m\eta+u_1-u_2)}{\sinh(m\eta)}X^2_{m-1}(u_2)Y^1_m(u_1),\end{aligned} \tag{A.3}$$

$$\begin{aligned}R_{1,2}(u_1-u_2)Y^1_m(u_1)X^2_{m+1}(u_2)&=\sinh(u_1-u_2)X^2_m(u_2)Y^1_{m-1}(u_1)\\&+\frac{\sinh\eta\sinh(m\eta-u_1+u_2)}{\sinh(m\eta)}Y^2_m(u_2)X^1_{m+1}(u_1),\end{aligned} \tag{A.4}$$

$$\overline{X}^1_{m+1}(u_1)\overline{X}^2_m(u_2)R_{1,2}(u_1-u_2)=\sinh(u_1-u_2+\eta)\overline{X}^2_{m+1}(u_2)\overline{X}^1_m(u_1), \tag{A.5}$$

$$\overline{Y}^1_{m-1}(u_1)\overline{Y}^2_m(u_2)R_{1,2}(u_1-u_2)=\sinh(u_1-u_2+\eta)\overline{Y}^2_{m-1}(u_2)\overline{Y}^1_m(u_1), \tag{A.6}$$

$$\begin{aligned}\overline{X}^1_{m-1}(u_1)\overline{Y}^2_m(u_2)R_{1,2}(u_1-u_2)&=\sinh(u_u-u_2)\overline{Y}^2_{m+1}(u_2)\overline{X}^1_m(u_1)\\&+\frac{\sinh\eta\sinh(m\eta+u_1-u_2)}{\sinh(m\eta)}\overline{X}^2_{m-1}(u_2)\overline{Y}^1_m(u_1),\end{aligned} \tag{A.7}$$

$$\begin{aligned}\overline{Y}^1_m(u_1)\overline{X}^2_{m+1}(u_2)R_{1,2}(u_1-u_2)&=\sinh(u_1-u_2)\overline{X}^2_m(u_2)\overline{Y}^1_{m-1}(u_1)\\&+\frac{\sinh\eta\sinh(m\eta-u_1+u_2)}{\sinh(m\eta)}\overline{Y}^2_m(u_2)\overline{X}^1_{m+1}(u_1).\end{aligned} \tag{A.8}$$

Multiplying Eq.(2.5) with $\overline{X}^1_{m'+1}(u_1)\overline{X}^2_{m'}(u_2)$ from the left and $X^1_m(u_1)X^2_{m-1}(u_2)$ from the right and using the relations (A.1) and (A.5), we arrive at (3.16). Multiplying Eq.(2.5) with $\overline{Y}^1_{m'+1}(u_1)\overline{X}^2_{m'+2}(u_2)$ from the left and $X^1_m(u_1)X^2_{m-1}(u_2)$ from the right and using the relations (A.1) and (A.8), we arrive at (3.17). Multiplying Eq.(2.5) with $\overline{X}^1_{m'+1}(u_1)\overline{X}^2_{m'}(u_2)$

from the left and $X^1_{m-2}(u_1)Y^2_{m-1}(u_2)$ from the right and using the relations (A.3) and (A.5), we arrive at (3.18) by exchanging u_1 and u_2.

Appendix B: Another gauge transformation

An alternating set of gauge matrices are

$$\widetilde{M}_k(u) = \left(\begin{array}{cc} \widetilde{X}_k(u), & \widetilde{Y}_k(u) \end{array} \right) = \left(\begin{array}{cc} \frac{e^{u-k\eta}}{\sinh(k\eta)} & e^{u+k\eta} \\ \frac{1}{\sinh(k\eta)} & 1 \end{array} \right), \tag{B.1}$$

$$\widetilde{M}^{-1}_k(u) = \left(\begin{array}{c} \widehat{Y}_k(u) \\ \widehat{X}_k(u) \end{array} \right) = \frac{e^{-u}}{2} \left(\begin{array}{cc} -1 & e^{u+k\eta} \\ \frac{1}{\sinh(k\eta)} & -\frac{e^{u-k\eta}}{\sinh(k\eta)} \end{array} \right). \tag{B.2}$$

Similarly, we define the following transformed matrices

$$\widetilde{L}^{(n)}_{j,k}(u) = \widetilde{M}^{-1}_j(u) L_n(u) \widetilde{M}_k(u), \tag{B.3}$$

$$\widetilde{T}_{j,k}(u) = \widetilde{M}^{-1}_j(u) T(u) \widetilde{M}_k(u) = \left(\begin{array}{cc} \widetilde{A}_{j,k}(u) & \widetilde{B}_{j,k}(u) \\ \widetilde{C}_{j,k}(u) & \widetilde{D}_{j,k}(u) \end{array} \right). \tag{B.4}$$

Following the similar procedures introduced in Section 3 and Section 4, we can construct the exact solutions for the case of $Re(\eta) < 0$.

References

[1] M. Toda, *J. Phys. Soc. Japan* **23**, 501 (1967)

[2] M. Toda, *Theory of Nonlinear Lattices*, Springer, 1981.

[3] M. Henon, *Phys. Rev.* **B 9** (1974), 1921.

[4] M. A. Olshanetsky and A. M. Perelomov, *Lett. Math. Phys.* **2** (1977), 7.

[5] H. Flaschka, *Prog. Theor. Phys.* **51** (1974), 703.

[6] A. Marshakov, *Seiberg-Witten Theory and Integrable Systems*, World Scientific, 1999.

[7] A. V. Mikhailov, M. A. Olshanetsky and A. M. Perelomov, *Commun. Math. Phys.* **79** (1981), 473.

[8] H. Flaschka, *Phys. Rev.* **B 9** (1974), 1924.

[9] H. Flaschka and D. W. McLaughlin, *Prog. Theor. Phys.* **55** (1976), 438.

[10] M. C. Gutzwiller, *Ann. Phys.* **133** (1981), 304.

[11] B. Sutherland, *Rocky Mountain J. Math.* **8** (1978), 413.

[12] E. K. Sklyanin, *Lect. Notes in Phys.* **226** (1985), 196.

[13] V. Pasquier and M. Gaudin, *J. Phys.* **A 25** (1992), 5243.

[14] E. K. Sklyanin, *J. Phys.* **A 46** (2013), 382001.

[15] M. Jimbo, *Comm. Math. Phys.* **102** (1986), 537.

[16] S. N. M. Ruijsenaars, *Comm. Math. Phys.* **133** (1990), 217.

[17] Yu. B. Suris, *Phys. Lett.* **A 145**, (1990), 113.

[18] A. Kundu, *Phys. Lett.* **A 190** (1994), 79.

[19] S. Pakuliak and S. Sergeev, *Czech. J. Phys.* **51** (2001), 1414.

[20] S. Kharchev and D. Lebedev, *Lett. Math. Phys.* **50** (1999), 53.

[21] F. A. Smirnov, *J. Phys.* **A 32** (1999), 5823.

[22] A. Gerasimov, S. Kharchev and D. Lebedev, *Int. Math. Res. Notices* **17** (2004), 823.

[23] L. A. Takhtadzhan and L. D. Faddeev, *Rush. Math. Surveys* **34** (1979), 11.

[24] J. Cao, H.-Q. Lin, K.-J. Shi and Y. Wang, *Nucl. Phys.* **B 663** (2003), 487.

[25] R. J. Baxter, *Exactly Solved Models in Statistical Mechanics*, Academic Press, 1982.

[26] V. E. Korepin, N. M. Bogoliubov and A. G. Izergin, *Quantum Inverse Scattering Method and Correlation Function*, Cambridge University Press, 1993.

[27] Y. Wang, W.-L. Yang, J. Cao and K. Shi, *Off-Diagonal Bethe Ansatz for Exactly Solvable Models*, Springer Press, 2015.

[28] X. Zhang, Y.-Y. Li, J. Cao, W.-L. Yang, K. Shi and Y. Wang, *Nucl. Phys.* **B 893** (2015), 70.

[29] X. Xu, J. Cao, S. Cui, W.-L. Yang, K. Shi and Y. Wang, *JHEP* **09** (2015), 212.